

<b>PATIENT INFORMATION</b> <b>Ropinirole (roe pin' i role") Extended-Release Tablets</b>	
<b>Important Note:</b> Ropinirole extended-release tablets have not been studied in RLS and is not approved for the treatment of RLS.	
<b>What is the most important information I should know about ropinirole extended-release tablets?</b> Ropinirole extended-release tablets can cause serious side effects including: <ul style="list-style-type: none"> <li>• <b>Falling asleep during normal activities.</b> You may fall asleep while doing normal activities such as driving a car, doing physical tasks, or using hazardous machinery while taking ropinirole extended-release tablets. You may suddenly fall asleep without being drowsy or without warning. This may result in having accidents. Your chances of falling asleep while doing normal activities while taking ropinirole extended-release tablets are greater if you take other medicines that cause drowsiness. Tell your healthcare provider right away if this happens. Before starting ropinirole extended-release tablets, be sure to tell your healthcare provider if you take any medicines that make you drowsy.</li> <li>• <b>Fainting.</b> Fainting can happen, and sometimes your heart rate may be decreased. This can happen especially when you start taking ropinirole extended-release tablets or your dose is increased. Tell your healthcare provider if you faint, feel dizzy or feel light-headed.</li> <li>• <b>Decrease in blood pressure.</b> Ropinirole extended-release tablets can decrease your blood pressure (hypotension), especially when you start taking ropinirole extended-release tablets or when your dose is changed. If you faint or feel dizzy, nauseated, or sweaty when you stand up from sitting or lying down (orthostatic hypotension), this may mean that your blood pressure is decreased. When you change position from lying down or sitting to standing up, you should do it carefully and slowly. Call your healthcare provider if you have any of the symptoms of decreased blood pressure listed above.</li> <li>• <b>Increase in blood pressure.</b> Ropinirole extended-release tablets may increase your blood pressure.</li> <li>• <b>Changes in heart rate (decrease or increase).</b> Ropinirole extended-release tablets can decrease or increase your heart rate.</li> <li>• <b>Hallucinations and other psychotic-like behavior.</b> Ropinirole extended-release tablets can cause or worsen psychotic-like behavior including hallucinations (seeing or hearing things that are not real), confusion, excessive suspicion, aggressive behavior, agitation, delusional beliefs (believing things that are not real), and disorganized thinking. The chances of having hallucinations or these other psychotic-like changes are higher in people with Parkinson's disease who are taking ropinirole extended-release tablets or taking higher doses of these drugs. If you have hallucinations or any of these other psychotic-like changes, talk with your healthcare provider.</li> <li>• <b>Uncontrolled sudden movements.</b> Ropinirole extended-release tablets may cause uncontrolled sudden movements or make such movements you already have worse or more frequent. Tell your healthcare provider if this happens. The doses of your anti-Parkinson's medicine may need to be changed.</li> <li>• <b>Unusual urges.</b> Some patients taking ropinirole extended-release tablets get urges to behave in a way unusual for them. Examples of this are an unusual urge to gamble, increased sexual urges and behaviors, or an uncontrollable urge to shop, spend money, or eat. If you notice or your family notices that you are developing any unusual behaviors, talk to your healthcare provider.</li> <li>• <b>Withdrawal Symptoms.</b> Ropinirole extended-release tablets are dopamine agonist medicines. Dopamine agonist medicines, including ropinirole extended-release tablets, can cause withdrawal symptoms as your dose is slowly lowered (tapered) or when treatment with ropinirole extended-release tablets is stopped. Tell your doctor right away if you get any of the following withdrawal symptoms:               <ul style="list-style-type: none"> <li>○ fever</li> <li>○ confusion</li> <li>○ severe muscle stiffness</li> <li>○ insomnia</li> <li>○ depression</li> <li>○ anxiety</li> <li>○ fatigue</li> <li>○ sweating</li> <li>○ pain</li> <li>○ feeling like you do not care about things you usually care about (apathy)</li> </ul> </li> </ul> After you have stopped taking ropinirole extended-release tablets, your healthcare provider may need to restart you at a low dose of ropinirole extended-release tablets if you get severe withdrawal symptoms.	
<b>What are ropinirole extended-release tablets?</b> <ul style="list-style-type: none"> <li>• Ropinirole extended-release tablets are long-acting prescription medicine containing ropinirole (taken 1 time a day) that is used only to treat Parkinson's disease but not to treat RLS.</li> </ul> Having one of these conditions does not mean you have or will develop the other condition. You should not be taking more than 1 medicine containing ropinirole. Tell your healthcare provider if you are taking any other medicine containing ropinirole. It is not known if ropinirole extended-release tablets are safe and effective for use in children younger than 18 years of age.	
<b>Do not take ropinirole extended-release tablets if you:</b> <ul style="list-style-type: none"> <li>• are allergic to ropinirole or any of the ingredients in ropinirole extended-release tablets. See the end of this page for a complete list of the ingredients in ropinirole extended-release tablets.</li> <li>• Get help right away if any of the symptoms of an allergic reaction cause problems swallowing or breathing.              Call your healthcare provider if you have any of the symptoms of an allergic reaction. Symptoms of an allergic reaction may include:               <ul style="list-style-type: none"> <li>○ hives</li> <li>○ swelling of the face, lips, mouth, tongue, or throat</li> <li>○ rash</li> <li>○ itching</li> </ul> </li> </ul>	
<b>Before taking ropinirole extended-release tablets, tell your healthcare provider about all of your medical conditions, including if you.</b> <ul style="list-style-type: none"> <li>• have daytime sleepiness from a sleep disorder to have unexpected or unpredictable sleepiness or periods of sleep.</li> <li>• start or stop taking other medicines while you are taking ropinirole extended-release tablets. This may increase your chances of getting side effects.</li> <li>• start or stop smoking while you are taking ropinirole extended-release tablets. Smoking may decrease the treatment effect of ropinirole extended-release tablets.</li> <li>• feel dizzy, nauseated, sweaty, or faint when you stand up from sitting or lying down.</li> <li>• drink alcoholic beverages. This may increase your chances of becoming drowsy or sleepy while taking ropinirole extended-release tablets.</li> <li>• have high or low blood pressure.</li> <li>• have or have had heart problems.</li> <li>• are pregnant or plan to become pregnant. It is not known if ropinirole extended-release tablets can harm your unborn baby.</li> <li>• are breastfeeding or plan to breastfeed. It is now known if ropinirole extended-release passes into your breast milk. The amount of breast milk you make may be decreased while taking ropinirole extended-release tablets. Talk to your healthcare provider to decide if you should breastfeed while taking ropinirole extended-release tablets.</li> </ul> Tell your healthcare provider about all the medicines you take, including prescription and over the counter medicines, vitamins, and herbal supplements. Some of these medicines may increase your chances of getting side effects while taking ropinirole extended-release tablets.	
<b>How should I take ropinirole extended-release tablets?</b> <ul style="list-style-type: none"> <li>• Take ropinirole extended-release tablets exactly as directed by your healthcare provider.</li> </ul>	

